



PHYSICS ACADEMY








CAREER SPECTRA

Institute for IIT-JAM | CSIR-NET/JRF | U-SET | GATE | JEST | TIFR | BARC

Dedicated to excellence...

GATE
CLASSICAL MECHANICS

PREVIOUS YEAR QUESTIONS WITH ANSWER
(CHAPTER-WISE)

-  **Newtonian Mechanics**
-  **Central Force**
-  **Special Theory of Relativity**
-  **Lagrangian**
-  **Hamiltonian**
-  **Poisson Bracket & Canonical Transformation**
-  **Phase Space Trajectory**

NEWTONIAN MECHANICS


- A particle is placed in a region with the potential $V(x) = \frac{1}{2} kx^2 - \frac{\lambda}{3} x^3$, where $k, \lambda > 0$.
Then, [GATE- 2010]

(a) $x=0$ and $x = \frac{k}{\lambda}$ are points of stable equilibrium
 (b) $x=0$ is a point of stable equilibrium and $x = \frac{k}{\lambda}$ is a point of unstable equilibrium
 (c) $x=0$ and $x = \frac{k}{\lambda}$ are points of unstable equilibrium
 (d) There are no points of stable or unstable equilibrium
- Two bodies of mass m and $2m$ are connected by a spring constant k . The frequency of the normal mode is. [GATE- 2011]

(a) $\sqrt{3k/2m}$ (b) $\sqrt{k/m}$ (c) $\sqrt{2k/3m}$ (d) $\sqrt{k/2m}$
- A particle of unit mass moves along the x -axis under the influence of a potential, $V(x) = x(x-2)^2$. The particle is found to be in stable equilibrium at the point $x=2$. The time period of oscillation of the particle is. [GATE- 2012]

(a) $\frac{\pi}{2}$ (b) π (c) $\frac{3\pi}{2}$ (d) 2π
- In the most general case, which one of the following quantities is NOT a second order tensor? [GATE- 2013]


(a) Stress (b) Strain
 (c) Moment of inertia (d) Pressure
- Consider two small blocks, each of mass M , attached to two identical springs. One of the springs is attached to the wall, as shown in the figure. The spring constant of each spring is k . The masses slide along the surface and the friction is negligible. The frequency of one of the normal modes of the system is, [GATE- 2013]



- (a) $\sqrt{\frac{3+\sqrt{2}}{2}} \sqrt{\frac{k}{M}}$ (b) $\sqrt{\frac{3+\sqrt{3}}{2}} \sqrt{\frac{k}{M}}$
 (c) $\sqrt{\frac{3+\sqrt{5}}{2}} \sqrt{\frac{k}{M}}$ (d) $\sqrt{\frac{3+\sqrt{6}}{2}} \sqrt{\frac{k}{M}}$

6. Two masses m and $3m$ are attached to the two ends of a massless spring with force constant K . If $m=100g$ and $K=0.3N/m$, then the natural angular frequency of oscillation is _____ Hz. [GATE- 2014]

7. A bead of mass m can slide without friction along a massless rod kept at 45° with the vertical as shown in the figure. The rod is rotating about the vertical axis with a constant angular speed ω . At any instant r is the distance of the bead from the origin. The momentum conjugate to r is: [GATE- 2014]



- (a) $m\dot{r}$ (b) $\frac{1}{\sqrt{2}} m\dot{r}$ (c) $\frac{1}{2} m\dot{r}$ (d) $\sqrt{2}m\dot{r}$

8. A particle of mass m is in a potential given by. [GATE- 2014]

$$V(r) = -\frac{a}{r} + \frac{ar_0^2}{3r^3}$$

Where a and r_0 are positive constants. When disturbed slightly from its stable equilibrium position it undergoes a simple harmonic oscillation. The time period of oscillation is

- (a) $2\pi\sqrt{\frac{mr_0^3}{2a}}$ (b) $2\pi\sqrt{\frac{mr_0^3}{a}}$
 (c) $2\pi\sqrt{\frac{2mr_0^3}{a}}$ (d) $4\pi\sqrt{\frac{mr_0^3}{a}}$

9. A particle of mass 0.01 kg falls freely in the earth's gravitational field with an initial Velocity $(0)=10\text{ms}^{-1}$. If the air exerts a frictional force of the form, $f = -kv$, then for $K=0.05\text{Nm}^{-1}\text{s}$, the velocity (in ms^{-1}) at time $t = 0.2\text{ s}$ is _____ (upto two decimal places). (use $g = 10\text{ms}^{-2}$ and $e = 2.72$): [GATE- 2015]

10. Consider the motion of the Sun with respect to the rotation of the Earth about its axis. If \vec{F}_c and \vec{F}_{co} denote the centrifugal and the Coriolis forces, respectively, acting on the Sun, then [GATE- 2015]


- (a) \vec{F}_c is radially outwards and $\vec{F}_{co} = \vec{F}_c$
 (b) \vec{F}_c is radially inwards and $\vec{F}_{co} = -2\vec{F}_c$
 (c) \vec{F}_c is radially outwards and $\vec{F}_{co} = -2\vec{F}_c$
 (d) \vec{F}_c is radially outwards and $\vec{F}_{co} = 2\vec{F}_c$

11. Two identical masses of 10 gm each are connected by a mass less spring of spring constant 1N/M. The non-zero angular eigen frequency of the system is.....rad/s. [GATE- 2017]
(up to two decimal places)
12. A uniform solid cylinder is released on a horizontal surface with speed 5m/ s without any rotation (slipping without rolling). The cylinder eventually starts rolling without slipping. If the mass and radius of the cylinder are 10 gm and 1cm respectively, the final linear velocity of the cylinder is..... m/ s . (up to two decimal places). [GATE- 2017]
13. A person weighs w_p at Earth's north pole and w_e at the equator. Treating the Earth as a perfect sphere of radius 6400 km , the value $100 \times \frac{(w_p - w_e)}{w_p}$ is.....(up to two decimal places). (Take $g = 10\text{ms}^{-2}$). [GATE- 2017]
14. In the context of small oscillations, which one of the following does NOT apply to the normal coordinates? [GATE- 2018]
(a) Each normal coordinate has an eigen-frequency associated with it
(b) The normal coordinates are orthogonal to one another
(c) The normal coordinates are all independent
(d) The potential energy of the system is a sum of squares of the normal coordinates with constant coefficients
15. The potential energy of a particle of mass m is given by. [GATE- 2020]
 $U(x) = \alpha \sin(k^2x - \pi/2), \quad \alpha > 0, k^2 > 0$
The angular frequency of small oscillations of the particle about $x = 0$ is
(a) $k^2 \sqrt{\frac{a}{m}}$ (b) $k^2 \sqrt{\frac{a}{2m}}$ (c) $k^2 \sqrt{\frac{2a}{m}}$ (d) $2k^2 \sqrt{\frac{a}{m}}$

CENTRAL FORCE

1. In a central force field, the trajectory of a particle of mass m and angular momentum L in plane polar coordinates is given by, [GATE- 2012]
$$\frac{1}{r} = \frac{m}{l^2}(1 + \epsilon \cos\theta)$$


Where, ϵ is the eccentricity of the particle's motion. Which one of the following choice for ϵ gives rise to a parabolic trajectory?
(a) $\epsilon = 0$ (b) $\epsilon = 1$ (c) $0 < \epsilon < 1$ (d) $\epsilon > 1$

2. A planet of mass m moves in a circular orbit of radius r in the gravitational potential $V(r) = -\frac{k}{r}$, where k is a positive constant. The orbit angular momentum of the planet is: [GATE- 2014]
 (a) $2r_0km$ (b) $\sqrt{2r_0km}$ (c) r_0km (d) $\sqrt{r_0km}$
3. A satellite is moving in a circular orbit around the Earth. If T, V and E are its average kinetic, average potential and total energies, respectively, then which one of the following options is correct? [GATE- 2015]
 (a) $V = -2T; E = -T$ (b) $V = -T; E = 0$
 (c) $V = -\frac{T}{2}; E = \frac{T}{2}$ (d) $V = \frac{-3T}{2}; E = \frac{-T}{2}$
4. An interstellar object has speed v at the point of its shortest distance R from a star of much larger mass M . Given $v^2 = 2GM/R$, the trajectory of the object is: [GATE- 2018]
 (a) Circle (b) Ellipse
 (c) Parabola (d) Hyperbola
5. A uniform circular disc of mass m and radius R is rotating with angular speed ω about an axis passing through its centre and making an angle $\theta = 30^\circ$ with the axis of the disc. If the kinetic energy of the disc is $\alpha m\omega^2 R^2$, the value of α is _____ (up to two decimal places). [GATE- 2018]
- 
6. A projectile of mass 1kg is launched at an angle of 30° from the horizontal direction at $t = 0$ and takes time T before hitting the ground. If its initial speed is 10ms^{-1} , the value of the action integral for the entire flight in the units of $\text{kgm}^2\text{s}^{-1}$ (round off to one decimal place) is _____. [Take $g = 10\text{ms}^{-2}$]. [GATE- 2019]
7. A particle is moving in a central force field given by $\vec{F} = -\frac{k}{r^3}\hat{r}$, where \hat{r} is the unit vector pointing away from the center of field. The potential energy of the particle is given by: [GATE- 2020]
 (a) $-\frac{K}{2r^2}$ (b) $-\frac{K}{r^2}$ (c) $\frac{K}{r^2}$ (d) $\frac{K}{2r^2}$


SPECIAL THEORY OF RELATIVITY

- For the set of all Lorentz transformations with velocities along the x -axis consider the two statements given below: [GATE- 2010]
 P: If L is a Lorentz transformation, then, L^{-1} is also a Lorentz transformation.
 Q: If L_1 and L_2 are Lorentz transformations, then $L_1 L_2$ is necessarily a Lorentz transformation.
 Choose the correct option
 (a) P is true and Q is false (b) Both P and Q are true
 (c) Both P and Q are false (d) P is false and Q is true
- A π^0 meson at rest decays into two photons, which moves along the x -axis. They are both detected simultaneously after a time, $t=10s$. In an inertial frame moving with a velocity $v=0.6c$ in the direction of one of the photons, the time interval between the two detections is. [GATE- 2010]
 (a) $15c$ (b) $0 s$ (c) $10 s$ (d) $20 s$
- Two particles each of rest mass m collide head-on and stick together. Before collision, the speed of each mass was 0.6 times the speed of light in free space. The mass of the final entity is. [GATE- 2011]
 (a) $5m/4$ (b) $2m$ (c) $5m/2$ (d) $25m/8$
- A rod of proper length l_0 oriented parallel to the x -axis moves with speed $2c / 3$ along the x -axis in the S -frame, where c is the speed of light in free space. The observer is also moving along the x -axis with speed $c / 2$ with respect to the S -frame. The length of the rod as measured by the observer is. [GATE- 2012]
 (a) $0.35l_0$ (b) $0.48l_0$ (c) $0.87l_0$ (d) $0.97l_0$
- An electron is moving with a velocity of $0.85c$ in the same direction as that of a moving photon. The relative velocity of the electron with respect to photon is: [GATE- 2013]
 (a) c (b) $-c$ (c) $0.15c$ (d) $-0.15c$
- The relativistic form of Newton's second law of motion is: [GATE- 2013]
 (a) $F = \frac{mc}{\sqrt{c^2-v^2}} \frac{dv}{dt}$ (b) $F = \frac{\sqrt{c^2-v^2}}{c} \frac{dv}{dt}$
 (c) $F = \frac{mc^2}{c^2-v^2} \frac{dv}{dt}$ (d) $F = \frac{c^2-v^2}{c^2} \frac{dv}{dt}$

7. If the half-life of an elementary particle moving with speed $0.9c$ in the laboratory frame is 5×10^{-8} s, then the proper half-life is _____ $\times 10^{-8}$ s. ($c = 3 \times 10^8$ m/s) [GATE- 2014]
8. In an inertial frame S , two events A and B take place at $(ct_A = 0, \vec{r}_A = 0)$ and $(ct_B = 0, \vec{r}_B = 2\hat{y})$, respectively. The times at which these events take place in a frame S' moving with a velocity $0.6c \hat{y}$ with respect to S are given by. [GATE- 2015]
 (a) $ct'_A = 0; ct'_B = -\frac{3}{2}$ (b) $ct'_A = 0; ct'_B = 0$
 (c) $ct'_A = 0; ct'_B = \frac{3}{2}$ (d) $ct'_A = 0; ct'_B = \frac{1}{2}$
9. A particle with rest mass M is at rest and decays into two particles of equal rest masses $\frac{3}{10}M$ which move along the z axis. Their velocities are given by: [GATE- 2015]
 (a) $\vec{v}_1 = \vec{v}_2 = (0.8c)\hat{z}$ (b) $\vec{v}_1 = -\vec{v}_2 = (0.8c)\hat{z}$
 (c) $\vec{v}_1 = -\vec{v}_2 = (0.6c)\hat{z}$ (d) $\vec{v}_1 = (0.6c)\hat{z}; \vec{v}_2 = (-0.8c)\hat{z}$
10. The kinetic energy of a particle of rest mass m_0 is equal to its rest mass energy. Its momentum in units of m_0c , where c is the speed of light in vacuum, is _____. [GATE- 2016]
 (Give your answer upto two decimal places)
11. In an inertial frame of reference S , an observer finds two events occurring at the same time at coordinates $x_1 = 0$ and $x_2 = d$. A different inertial frame S' moves with velocity v with respect to S along the positive x -axis. An observer in S' also notices these two events and finds them to occur at times t'_1 and t'_2 and at positions x'_1 and x'_2 respectively. [GATE- 2016]
 If $\Delta t' = t'_2 - t'_1, \Delta x' = x'_2 - x'_1$ and $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$, which of the following statement is true?
 (a) $\Delta t' = 0, \Delta x' = \gamma d$ (b) $\Delta t' = 0, \Delta x' = \frac{d}{\gamma}$
 (c) $\Delta t' = \frac{-\gamma v d}{c^2}, \Delta x' = \gamma d$ (d) $\Delta t' = \frac{-\gamma v d}{c^2}, \Delta x' = \frac{d}{\gamma}$
12. A particle of rest mass M is moving along the positive x -direction. It decays into two photons γ_1 and γ_2 as shown in the figure. The energy of γ_1 is 1 GeV and the energy of γ_2 is 0.82 GeV. The value of M (in units of $\frac{\text{GeV}}{c^2}$) is _____. [GATE- 2016]
 (Give your answer upto two decimal places)




13. An object travels along the x -direction with velocity $\frac{c}{2}$ in a frame O . An observer in a frame O' sees the same object travelling with velocity $\frac{c}{4}$. The relative velocity of O' with respect to O in units of c is(up to two decimal places). [GATE- 2017]
14. A spaceship is travelling with a velocity of $0.7c$ away from a space station. The spaceship ejects a probe with a velocity $0.59c$ opposite to its own velocity. A person in the space station would see the probe moving at a speed Xc , where the value of X (up to three decimal places). [GATE- 2018]
15. Two spaceships A and B , each of the same rest length L , are moving in the same direction with speeds $\frac{4c}{5}$ and $\frac{3c}{5}$, respectively, where c is the speed of light. As measured by B , the time taken by A to completely overtake B [see figure below] in units of L/c (to the nearest integer) is _____



16. Two events, one on the earth and the other one on the Sun, occur simultaneously in the earth's frame. The time difference between the two events as seen by an observer in a spaceship moving with velocity $0.5c$ in the earth's frame along the line joining the earth to the Sun is Δt , where c is the speed of light. Given that light travels from the Sun to the earth in 8.3 minutes in the earth's frame, the value of $|\Delta t|$ in minutes (rounded off to two decimal places) is _____ (Take the earth's frame to be inertial and neglect the relative motion between the earth and the sun). [GATE- 2019]

LAGRANGIAN

1. A particle is moving under the action of a generalized potential $V(q, \dot{q}) = \frac{1+\dot{q}}{q^2}$. The magnitude of the generalized force is: [GATE- 2011]
 (a) $\frac{2(1+\dot{q})}{q^3}$ (b) $\frac{2(1-\dot{q})}{q^3}$ (c) $\frac{2}{q^3}$ (d) $\frac{\dot{q}}{q^3}$
2. A particle of mass m slides under the gravity without friction along the parabolic path $y=ax^2$, as shown in the figure. Here a is a constant. [GATE- 2012]



The Lagrangian for this particle is given by

- (a) $L = \frac{1}{2} m \dot{x}^2 - mgax^2$
 - (b) $L = \frac{1}{2} m(1+4a^2x^2)\dot{x}^2 - mgax^2$
 - (c) $L = \frac{1}{2} m \dot{x}^2 + mgax^2$
 - (d) $L = \frac{1}{2} m(1+4a^2x^2)\dot{x}^2 + mgax^2$
3. The Lagrange's equation of motion of the particle for above question is given by
 (a) $\ddot{x} = 2gax$
 (b) $m(1+4a^2x^2)\ddot{x} = -2mgax - 4ma^2x\dot{x}^2$
 (c) $m(1+4a^2x^2)\ddot{x} = 2mgax + 4ma^2x\dot{x}^2$
 (d) $\ddot{x} = -2gax$
4. The Lagrangian of a system with one degree of freedom q is given by $L = a\dot{q}^2 + \beta q^2$, where α and β are non-zero constants. If p_q denotes the canonical momentum conjugate to q then which one of the following statements is CORRECT? [GATE- 2013]
 (a) $p_q = 2\beta q$ and it is a conserved quantity.
 (b) $p_q = 2\beta q$ and it is not a conserved quantity.
 (c) $p_q = 2\alpha \dot{q}$ and it is a conserved quantity.
 (d) $p_q = 2\alpha \dot{q}$ and it is not a conserved quantity.
5. The Lagrangian for a particle of mass m at a position \vec{r} moving with a velocity \vec{v} is given by $L = \frac{m}{2} \vec{v}^2 + C\vec{r} \cdot \vec{v} - V(r)$, where $V(r)$ is a potential and C is a constant. If \vec{p}_c is the canonical momentum, then its Hamiltonian is given by: [GATE- 2015]
 (a) $\frac{1}{2m}(\vec{p}_c + C\vec{r})^2 + V(r)$ (b) $\frac{1}{2m}(\vec{p}_c + C\vec{r})^2 - V(r)$

(c) $\frac{p_c^2}{2m} + V(r)$

(d) $\frac{1}{2m} p_c^2 + C^2 r^2 + V(r)$

6. The Lagrangian of a system is given by. [GATE- 2016]

$L = \frac{1}{2}ml^2[\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2] - mgl \cos \theta$, where m, l and g are constants.

Which of the following is conserved?

- (a) $\dot{\phi} \sin^2 \theta$ (b) $\dot{\phi} \sin \theta$ (c) $\frac{\dot{\phi}}{\sin \theta}$ (d) $\frac{\dot{\phi}}{\sin^2 \theta}$

7. If the Lagrangian $L_0 = \frac{1}{2}m \left(\frac{dq}{dt}\right)^2 - \frac{1}{2}m\omega^2 q^2$ is modified to $L = L_0 + \alpha q \left(\frac{dq}{dt}\right)$, which one of the following is TRUE? [GATE- 2017]

- (a) Both the canonical momentum and equation of motion do not change
 (b) Canonical momentum changes, equation of motion does not change
 (c) Canonical momentum does not change, equation of motion changes
 (d) Both the canonical momentum and equation of motion change

8. Consider the Hamiltonian $H(q, p) = \frac{ap^2 q^4}{2} + \frac{\beta}{q^2}$, where α and β are parameters with appropriate dimensions, and q and p are the generalized coordinate and momentum, respectively. The corresponding Lagrangian $L(q, \dot{q})$ is: [GATE- 2019]

- (a) $\frac{1}{2\alpha} \frac{\dot{q}^2}{q^4} - \frac{\beta}{q^2}$ (b) $\frac{1}{2\alpha} \frac{\dot{q}^2}{q^4} + \frac{\beta}{q^2}$ (c) $\frac{1}{2\alpha} \frac{\dot{q}^2}{q^4} + \frac{\beta}{q^2}$ (d) $-\frac{1}{2\alpha} \frac{\dot{q}^2}{q^4} + \frac{\beta}{q^2}$

9. Consider the Lagrangian $L = a \left(\frac{dx}{dy}\right)^2 + b \left(\frac{dx}{dt}\right)^2 + cxy$, where a, b and c constants. If p_x and p_y are the momenta conjugate to the coordinates x and y respectively, then the Hamiltonian is. [GATE- 2020]

- (a) $\frac{p_x^2}{a} + \frac{p_y^2}{b} + cxy$ (b) $\frac{p_x^2}{2a} + \frac{p_y^2}{2b} - cxy$
 (c) $\frac{p_x^2}{4a} + \frac{p_y^2}{4b} - cxy$ (d) $\frac{p_x^2}{2a} + \frac{p_y^2}{2b} + cxy$


HAMILTONIAN

1. Hamilton's equations are then given by: [GATE- 2010]

- (a) $\dot{p}_\theta = -mgl \sin \theta; \dot{\theta} = \frac{p_\theta}{ml^2}$ (b) $\dot{p}_\theta = mgl \sin \theta; \dot{\theta} = \frac{p_\theta}{ml^2}$
 (c) $\dot{p}_\theta = -m\dot{\theta}; \dot{\theta} = \frac{p_\theta}{m}$ (d) $\dot{p}_\theta = -\left(\frac{g}{l}\right)\theta; \dot{\theta} = \frac{p_\theta}{ml}$

2. A particle of mass m is attached to a fixed point O by a weightless inextensible string of length a . It is rotating under the gravity as shown in the figure. The Lagrangian of the particle is. [GATE- 2012]

$L(\theta, \dot{\theta}, \phi, \dot{\phi}) = \frac{1}{2}ma^2(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) - mga \cos \theta$ where θ and ϕ are the polar angles. The Hamiltonian of the particles is



- (a) $H = \frac{1}{2ma^2} \left(p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} \right) - mga \cos \theta$ (b) $H = \frac{1}{2ma^2} \left(p_\theta^2 + \frac{p_\phi^2}{\sin^2 \theta} \right) + mga \cos \theta$
- (c) $H = \frac{1}{2ma^2} (p_\theta^2 + p_\phi^2) - mga \cos \theta$ (d) $H = \frac{1}{2ma^2} (p_\theta^2 + p_\phi^2) + mga \cos \theta$

3. The Hamilton's canonical equation of motion in terms of Poisson Brackets are [GATE- 2014]

- (a) $\dot{q} = \{q, H\}; \dot{p} = \{p, H\}$ (b) $\dot{q} = \{H, q\}; \dot{p} = \{H, p\}$
 (c) $\dot{q} = \{H, p\}; \dot{p} = \{H, q\}$ (d) $\dot{q} = \{p, H\}; \dot{p} = \{q, H\}$

4. The Hamiltonian for a system of two particles of masses m_1 and m_2 at \vec{r}_1 and \vec{r}_2 having velocities \vec{v}_1 and \vec{v}_2 is given by $H = \frac{1}{2} m_1 v_1^2 + \frac{1}{2} m_2 v_2^2 + \frac{C}{(\vec{r}_1 - \vec{r}_2)^2} \hat{z} \cdot (\vec{r}_1 \times \vec{r}_2)^2$, where C is constant. Which one of the following statements is correct? [GATE- 2015]

- (a) The total energy and total momentum are conserved
 (b) Only the total energy is conserved
 (c) The total energy and the z - component of the total angular momentum are conserved
 (d) The total energy and total angular momentum are conserved

5. If H is the Hamiltonian for a free particle with mass m , the commutator $[x, [x, H]]$ is. [GATE- 2018]

- (a) \hbar^2/m (b) $-\hbar^2/m$
 (c) $-\hbar^2/(2m)$ (d) $\hbar^2/(2m)$

6. The Hamiltonian for a particle of mass m is $H = \frac{p^2}{2m} + kqt$ where q and p are the generalized coordinate and momentum, respectively, t is time and k is a constant.

For the initial condition, $q = 0$ and $p = 0$ at $t = 0, q(t) \propto t^a$. The value of α is _____ . [GATE- 2019]

7. A small disc is suspended by a fiber such that it is free to rotate about the fiber axis (see figure). For small angular deflections, the Hamiltonian for the disc is given by $H = \frac{p_\theta^2}{2I} + \frac{1}{2} \alpha \theta^2$, Where I is the moment of inertia and α is the restoring torque per unit deflection. The disc is subjected to angular deflections (θ) due to thermal collisions from the surroundings gas at temperature T and p_θ is the momentum conjugate to θ . The average and the root-mean-square angular deflection, θ_{avg} and θ_{rms} , respectively are _____ [GATE- 2020]



- (a) $\theta_{avg} = 0$ and $\theta_{rms} = \left(\frac{k_B T}{\alpha}\right)^{1/2}$
- (b) $\theta_{avg} \neq 0$ and $\theta_{rms} = \left(\frac{k_B T}{\alpha}\right)^{1/2}$
- (c) $\theta_{avg} \neq 0$ and $\theta_{rms} = \left(\frac{k_B T}{\alpha}\right)^{3/2}$
- (d) $\theta_{avg} = 0$ and $\theta_{rms} = \left(\frac{k_B T}{\alpha}\right)^{3/2}$

POISSON BRACKET & CANONICAL TRANSFORMATION

1. The Poisson bracket between θ and $\dot{\theta}$ is: [GATE- 2010]
 - (a) $\{\theta, \dot{\theta}\} = 1$
 - (b) $\{\theta, \dot{\theta}\} = \frac{1}{ml^2}$
 - (c) $\{\theta, \dot{\theta}\} = \frac{1}{m}$
 - (d) $\{\theta, \dot{\theta}\} = \frac{g}{l}$
2. The Poisson bracket $[x, xp_y + yp_x]$ is equal to. [GATE- 2017]
 - (a) $-x$
 - (b) y
 - (c) $2p_x$
 - (d) p_y
3. Let (p, q) and (P, Q) be two pairs of canonical variables. The transformation $Q = q^\alpha \cos(\beta p), P = q^\alpha \sin(\beta p)$ is canonical for. [GATE- 2011]
 - (a) $a=2, \beta = \frac{1}{2}$
 - (b) $a=2, \beta=2$
 - (c) $a=1, \beta=1$
 - (d) $a=\frac{1}{2}, \beta=2$
4. Given that the linear transformation of a generalized coordinate q and the corresponding momentum p , $Q = q + 4ap, P = q + 2p$ is canonical, the value of the constant a is _____
5. For the transformation [GATE- 2018]

$$Q = \sqrt{2q} e^{-1+2a} \cos p, P = \sqrt{2q} e^{-a-1} \sin p$$
 (where α is a constant) to be canonical, the value of α is _____.

6. Consider a transformation from one set of generalized coordinate and momentum (q, p) to another set (Q, P) denoted by, [GATE- 201]


$$Q = pq^s; \quad P = q^r$$

Where s and r are constants. The transformation is canonical if

- (a) $s = 0$ and $r = 1$ (b) $s = 2$ and $r = -1$
 (c) $s = 0$ and $r = -1$ (d) $s = 2$ and $r = 1$
7. Let p be the momentum conjugate to the generalized coordinate q . If the transformation [GATE- 2020]
- $$Q = \sqrt{2}q^m \cos p$$
- $$P = \sqrt{2}q^m \sin p$$
- Is canonical, then $m = \underline{\hspace{2cm}}$.
8. Let u^μ denote the 4-velocity of a relativistic particle whose square $u^\mu u_\mu = 1$. If $\epsilon_{\mu\nu\rho\sigma}$ is the Levi-Civita tensor then the value of $\epsilon_{\mu\nu\rho\sigma} u^\mu u^\nu u^\rho u^\sigma$ is $\underline{\hspace{2cm}}$. [GATE- 2020]


PHASE SPACE TRAJECTORY

1. The Hamiltonian of particle of mass m is given by $H = \frac{p^2}{2m} - \frac{aq^2}{2}$. Which one of the following figure describes the motion of the particle in phase space? [GATE- 2014]




2. The phase space trajectory of an otherwise free particle bouncing between two hard walls elastically in one dimension is a. [GATE- 2017]
- (a) straight line (b) parabola (c) rectangle (d) circle

3. A particle moves in one dimension under a potential $V(x) = \alpha |x|$ with some non-zero total energy. Which one of the following best describes the particle trajectory in the phase space? [GATE- 2018]



4. A ball bouncing of a rigid floor is described by the potential energy function $V(x) = \begin{cases} mgx & \text{for } x > 0 \\ \infty & \text{for } x \leq 0 \end{cases}$ [GATE- 2019]

Which of the following schematic diagrams best represents the phase space plot of the ball?



ANSWER-KEY

NEWTONIAN MECHANICS

1.	B	2.	A	3.	B	4.	B	5.	C	6.	0.318	7.	A
8.	A	9.	4.94	10.	B	11.	14.14	12.	3.33	13.	0.33	14.	B
15.	*	16.		17.		18.		19.		20.		21.	

CENTRAL FORCE

1.	B	2.	D	3.	S	4.	C	5.	0.21	6.	33.3	7.	*

SPECIAL THEORY OF RELATIVITY

1.	B	2.	A	3.	C	4.	D	5.	B	6.	C	7.	2.18
8.	A	9.	B	10.	1.73	11.	C	12.	1.44	13.	0.28	14.	0.187c
15.	5	16.	4.77										

LAGRANGIAN

1.	C	2.	B	3.	B	4.	D	5.	B	6.	A	7.	B
8.	A	9.	*										

HAMILTONIAN

1.	A	2.	B	3.	A	4.	C	5.	B	6.	3	7.	*
----	---	----	---	----	---	----	---	----	---	----	---	----	---

POSSION BRACKET & CANONICAL TRANSFORMATION

1.	B	2.	B	3.	D	4.	0.25	5.	2	6.	B	7.	0.5
8.	6												

PHASE SPACE TRAJECTORY

1.	D	2.	C	3.	A	4.	B						
----	---	----	---	----	---	----	---	--	--	--	--	--	--

